Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MECHANICS.

161. Proposed by **W. J. GREENSTREET, A. M.**, Editor of The Mathematical Gazette, Stroud, England.

Four equal uniform smoothly jointed rods length a, and weight w, form a rhombus $ABCD$, A and C being in contact with two vertical walls b feet apart. An elastic string, natural length x, modulus λ, keeps the figure in position. The angle of friction at A and C is $\tan^{-1}p$. When the rhombus is just about to slip, find the angle A, and the angle between AB and the vertical.

162. Proposed by **B. F. FINKEL, A.M., M.Sc.**, Professor of Mathematics and Physics, Drury College, Springfield, Mo.

Show that the velocity, v, of a wave along the surface of a liquid whose depth is not less than the wave-length, λ, whose density is δ, and surface tension, T, is $v^2=\dfrac{g\lambda}{2\pi}+\dfrac{2\pi T}{\lambda\delta}$.

DIOPHANTINE ANALYSIS.

118. Proposed by **L. C. WALKER, A.M.**, Professor of Mathematics, Colorado School of Mines, Golden, Col.

Find the two least integral numbers such that their sum shall be a square, and the sum of their squares a biquadrate.

AVERAGE AND PROBABILITY.

147. Proposed by **W. J. GREENSTREET, A. M.**, Editor of The Mathematical Gazette, Stroud; England.

In a bag are n balls, known to be black or white, either color, *a priori*, equally likely. I draw two, which turn out to be one white and one black. I replace them and draw two more. What is the chance both are black?

◆

NOTES.

Professor Lon C. Walker has been elected Professor of Mathematics in the Colorado School of Mines, Golden, Colorado.

Dr. Saul Epsteen, of the University of Chicago, has very kindly consented to edit, for the coming year, the problems and solutions for the MONTHLY. So beginning with this issue all problems and solutions should be sent to him. The MONTHLY is to be congratulated that it has fallen into the hands of such able mathematicians as Drs. Dickson and Epsteen. We predict for it the most prosperous year in its history. B. F. F.

ERRATA.

On page 171, problem 105, for "$a^{2m}+b^{2m}$" read $a^{2^m}+b^{2^m}$.